aldehyde. Vitamin A Wittig reagent and vitamin A aldehyde are reacted together to form *beta*-carotene.

(b) The ingredient meets the specifications of the Food Chemicals Codex, 3d Ed. (1981), p. 73, which is incorporated by reference. Copies are available from the National Academy Press, 2101 Constitution Ave. NW., Washingtion, DC 20418, or available for inspection at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030, or go to: http://www.archives.gov/federal register/

code_of_federal_regulations/ibr_locations.html.

- (c) In accordance with §184.1(b)(1), the ingredient is used in food with no limitation other than current good manufacturing practice. The affirmation of this ingredient as generally recognized as safe (GRAS) as a direct human food ingredient is based upon the following current good manufacturing practice conditions of use:
- (1) The ingredient is used as a nutrient supplement as defined in §170.3(o)(20) of this chapter.
- (2) The ingredient is used in the following foods at levels not to exceed current good manufacturing practice: dairy product analogs as defined in §170.3(n)(10) of this chapter; fats and oils as defined in §170.3(n)(12) of this chapter; and processed fruits and fruit juices as defined in §170.3(n)(35) of this chapter. Beta-carotene may be used in infant formula as a source of vitamin A in accordance with section 412(g) of the Federal Food, Drug, and Cosmetic Act or with regulations promulgated under section 412(g) of the act.
- (d) Prior sanctions for this ingredient different from the uses established in this section do not exist or have been waived.

 $[52~{\rm FR}~25211,~{\rm July}~6,~1987]$

§ 184.1250 Cellulase enzyme preparation derived from Trichoderma longibrachiatum.

(a) Cellulase enzyme preparation is derived from a nonpathogenic, nontoxicogenic strain of *Trichoderma longibrachiatum* (formerly *T. reesei*). The enzyme, cellulase, catalyzes the endohydrolysis of 1,4-beta-glycosidic

linkages in cellulose. It is obtained from the culture filtrate resulting from a pure culture fermentation process.

- (b) The ingredient meets the general and additional requirements for enzyme preparations in the monograph specifications on enzyme preparations in the "Food Chemicals Codex." 4th ed. (1996), pp. 129 to 134, which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies are available from the National Academy Press, 2101 Constitution Ave. NW., Box 285, Washington, DC 20055 (Internet http://www.nap.edu), or may be examined at the Center for Food Safety and Applied Nutrition's Library, 5100 Paint Branch Pkwy., College Park, MD 20740, or at the National Archives and Records Administration (NARA). For information on the availability of this material at NARA, call 202-741-6030, orgoto: http:// www.archives.gov/federal register/ code of federal regulations/ ibr locations.html.
- (c) In accordance with §184.1(b)(1), the ingredient is used in food with no limitation other than current good manufacturing practice. The affirmation of this ingredient as generally recognized as safe (GRAS) as a direct human food ingredient is based upon the following current good manufacturing practice conditions of use:
- (1) The ingredient is used in food as an enzyme as defined in §170.3(o)(9) of this chapter for the breakdown of cellulose.
- (2) The ingredient is used in food at levels not to exceed current good manufacturing practice.

[64 FR 28361, May 26, 1999]

§ 184.1257 Clove and its derivatives.

(a) Cloves are the dried unopened flower buds and calyx tubes, harvested before the flowers have opened, of the clove tree *Eugenia caryophyllata* Thunberg, native to tropical Asia. Their derivatives include essential oils (cloves, CAS Reg. No. 8005–34–8; buds; leaves, CAS Reg. No. 8015–97–2; stems, CAS Reg. No. 8015–98–3; and eugenol, CAS Reg. No. 97–53–0), oleoresins, and natural extractives obtained from clove buds, leaves, and stems.